



ELSEVIER

INTERNATIONAL SOCIETY
FOR INFECTIOUS DISEASES<http://intl.elsevierhealth.com/journals/ijid>

LETTER TO THE EDITOR

***Streptococcus pyogenes* bacteremia in the setting of cholangitis**

Streptococcus pyogenes bacteremia occurs most commonly in the setting of skin and wound infections, followed by respiratory tract infections.¹ *S. pyogenes* has never been reported to cause cholangitis. Herein we describe a case of cholangitis in the setting of *S. pyogenes* bacteremia.

A 52-year-old man underwent cholecystojejunostomy and gastrojejunostomy for pancreatic cancer in February 2002. His course was complicated by obstructive jaundice and *Escherichia coli* bacteremia in September 2002. He underwent percutaneous transhepatic cholangiography with biliary drainage and received ofloxacin for 14 days. An internal biliary stent was then inserted percutaneously. Two months later he was admitted with chills, abdominal pain and jaundice. He was afebrile but had icterus and hepatomegaly. He had no evidence of pharyngeal erythema, lung or skin infection. White blood cell count was $5.7 \times 10^9/L$ with 65% neutrophils. Serum bilirubin, alkaline phosphatase, and γ -glutamyl transpeptidase were markedly elevated compared to values obtained on the previous admission. Later, blood cultures grew group A β -hemolytic *Streptococcus* spp. Computerized tomography of the abdomen revealed intra- and extrahepatic biliary ductal dilatation. The patient was treated with ampicillin and underwent endoscopic biliary stent placement with marked clinical improvement. Since no other organisms were isolated from blood cultures, the patient was diagnosed with *S. pyogenes* cholangitis.

The site of origin of the infecting organisms in cholangitis remains debatable.² Gram-negative enteric bacilli and anaerobic bacteria are commonly incriminated and bacteremia occurs in 50% of patients. The most frequently isolated organism is *E. coli*, followed by *Bacteroides fragilis*.³ Bacteremia due to *S. pyogenes* is an uncommon condition. It has been described in the setting of erysipelas, intraabdominal infections or overwhelming septicemia,^{4–6} but has not been previously reported to occur with cholangitis. Our patient had clinical, laboratory and radiological findings highly suggestive of cholangitis. At the

time of the endoscopic retrograde cholangiography, obstruction of the old biliary stent was noted. The source of the streptococcal infection is not certain, but it could have been due to colonization of the biliary stent with *S. pyogenes*. Obstruction of the stent and the subsequent bile stagnation provided a favorable milieu for the growth of this organism, resulting in cholangitis and bacteremia.

Some references have pointed out that group A streptococci do not usually grow in bile,⁷ and this is used in vitro to differentiate these organisms from enteric streptococci. It should be noted, however, that this property is also shared by group C, G and viridans streptococci. We found three reports of cholangitis caused by viridans streptococci in the literature.^{8–10} This makes cholangitis due to *S. pyogenes* theoretically plausible. It could be that in cases of high inoculum, the bacteria are able to survive in bile and cause clinical disease. In this particular case, we did not test the *S. pyogenes* isolate recovered from blood for its ability to grow in bile.

Although enteric bacteria are the organisms most commonly incriminated in cholangitis, other organisms of non-enteric origin such as *S. pyogenes* may cause cholangitis, especially in the presence of a foreign body in the biliary tree.

Conflict of interest: No conflicting interest declared.

References

1. Duma RJ, Weinberg AN, Medrek TF, Kunz LJ. Streptococcal infections. A bacteriologic and clinical study of streptococcal bacteremia. *Medicine* 1969;48:87–127.
2. Sievert W, Vakil NB. Emergencies of the biliary tract. *Gastroenterol Clin North Am* 1988;17:245–64.
3. Levison ME, Bush LM. Peritonitis and other intra-abdominal infections. In: Mandell GL, Bennett JE, Dolin R, editors. *Mandell, Douglas, and Bennett's principles and practice of infectious diseases*. Philadelphia: Churchill Livingstone; 2000, p. 821–856.
4. Dan M, Maximova S, Siegman-Igra Y, Gutman R, Rotmensch HH. Varied presentations of sporadic group A streptococcal bacteremia: clinical experience and attempt at classification. *Rev Infect Dis* 1990;12:537–42.

5. Goepel JR, Richards DG, Harris DM, Henry L. Fulminant streptococcus pyogenes infection. *BMJ* 1980;**281**:1412.
6. Ispahani P, Donald FE, Aveline AJ. *Streptococcus pyogenes* bacteraemia: an old enemy subdued, but not defeated. *J Infect* 1988;**16**:37–46.
7. Tapsall JW, Chau C. Beta-haemolytic bacitracin sensitive strains of *Streptococcus salivarius*: a source of confusion with *Streptococcus pyogenes* in throat swab cultures. *Pathology* 1980;**12**:403–6.
8. Turner FP, Ohler RL. Suppurative cholangitis with multiple liver abscess formation due to *Aerobacter aerogenes* and *Streptococcus viridans*: Successful treatment with massive doses of intravenous penicillin. *J Maine Med Assoc* 1965;**56**:177–80.
9. Altucci P, Catalano G, Gattoni A, Onza I, Abbate GF. Isolation of protoplasts from the blood in a case of *Streptococcus viridans* cholangitis. *Boll Ist Sieroter Milan* 1972;**51**:183–6.
10. Murray HW, Gross KC, Masur H, Roberts RB. Serious infections caused by *Streptococcus milleri*. *Am J Med* 1978;**64**:759–64.

Ghenwa K. Dakdouki
Zeina A. Kanafani
Fadi H. Mourad
Ali I. Shamseddine
Souha S. Kanj*

*Department of Internal Medicine
American University of Beirut
Medical Center, P.O. Box 113-6044
Hamra 11032090 Beirut, Lebanon*

*Corresponding author

Tel.: +961-1-350000; fax: +961-1-370814

E-mail address: sk11@aub.edu.lb
(S.S. Kanj)

5 November 2003

Corresponding Editor:

Jonathan Cohen, Brighton, UK

Available online at www.sciencedirect.com

